Received 27 August 2015

(wileyonlinelibrary.com) DOI: 10.1002/mma.3908 MOS subject classification: 35C08; 35Q51; 65M06

Interaction of sine-Gordon solitons in the model with attracting impurities

Evgenii G. Ekomasov, Azamat M. Gumerov^{*†} and Ramil R. Murtazin

Communicated by Prof. Amar Debbouche

We study properties of the localized solutions to the sine-Gordon equation excited on the attractive impurity by a moving kink. The cases of one-dimensional and two-dimensional spatially extended impurities are considered. For the case of one-dimensional impurity the possibility of excitation of the first even and odd high-amplitude impurity modes by the moving kink is demonstrated. By linearizing the sine-Gordon equation the dispersion relations for the small-amplitude localized impurity modes were obtained. The numerically obtained dispersion relations in the case of low oscillation amplitudes are in a good agreement with the results of analytical calculations. For the case of two-dimensional impurity we show the possibility of excitation of the nonlinear high-amplitude waves of new type called here a breathing pulson and a breathing 2D soliton. We suggest analytical expressions to model these nonlinear excitations. The breathing pulson and breathing 2D soliton are long-lived and can be of both symmetric and asymmetric type depending on the impurity type. The range of the impurity parameters where the breathing pulson and breathing 2D soliton can be excited was determined. The dependencies of the oscillation frequency and the amplitude of the excited impurity modes on the impurity parameters are reported. Copyright © 2016 John Wiley & Sons, Ltd.

Keywords: sine-Gordon equation; impurity; kink; soliton; pulson

1. Introduction

Solitons, initially studied in integrable systems, gave rise to the study on solitary waves in non-integrable systems to describe a number of physical phenomena. For example, the sin-Gordon equation (SGE) is used for modeling wave propagation in geological media, in molecular biology, field theory models, in elementary particle physics, to name a few [1–3]. The SGE soliton solutions help to describe domain walls in magnetics, dislocations in crystals, fluxons in Josephson's junctions, etc. [4–6]. In many cases behavior of solitons can be effectively described by quasi-particles and then their dynamics can be presented by ordinary differential equations [7]. However, in the presence of perturbations, the structure of solitons can be changed and they can be more accurately described by deformable quasi-particles [5]. Soliton's internal degrees of freedom can be excited in this case and they can play a very important role in a number of physical processes [8–12]. Soliton internal modes can be responsible for the non-trivial effects of their interactions [13, 14]. The internal modes can include the translational and pulsation modes describing long-lived oscillations of soliton width [15]. The effect of various perturbations on the excitation of SGE soliton's internal modes attracts a lot of attention of researchers. Local inhomogeneities are ubiquitous in many physical systems, including those described by SGE, and it is very important to study the soliton scattering on such impurities [5]. For instance, there exist many works devoted to the analysis of external force that varies in time and space [5, 9, 15–17]. Weak perturbations on the SGE solutions can be studied in frame of the perturbation theory well-developed for solitons [5, 7] but the effect of strong perturbations is more often addressed numerically [18–21].

It is also very interesting to study the effect of an impurity in the system modeled by spatial modulation (inhomogeneity) of the periodic potential [5]. Depending on the geometry of the system, one-dimensional or multi-dimensional problems can be physically meaningful. The problem of scattering of SGE kinks on impurities in one-dimensional case has been under consideration for a long time [7, 19, 22, 23]. For example, the model of classical particle is applicable to the problem of kink-impurity interaction in the case when impurity itself does not support vibrational modes localized on the impurity [5]. The mechanisms of kink-impurity interactions has been demonstrated in the works [5, 24–27]. Let us mention such an interesting effect as the reflection of a kink by an attractive impurity due to the resonance energy exchange between translational kink's mode and the impurity mode. Two-dimensional SGE has been also studied for a long time with the use of analytical methods [28–32] and with the help of numerical methods [18, 28, 33]. For

Bashkir State University, Zaki Validi Str., 32, Ufa 450076, Russia

^{*} Correspondence to: Åzamat M. Gumerov, Bashkir State University, Zaki Validi Str., 32, Ufa 450076, Russia.

[†] E-mail: bgu@bk.ru

example, in the works [18, 34] the appearance and motion of flexural solitary waves on the kink interacting with a two-dimensional impurity has been investigated. However, the possibility of excitation of various two-dimensional vibrational modes localized on the impurity was not discussed in those works.

In the present study we analyze the interaction of SGE kink with an impurity in the case when large-amplitude nonlinear waves localized on the impurity are excited as a result of interaction.

Let us consider the system defined by the following Lagrangian

$$L(x, y, t) = \frac{1}{2} \left[\left(\frac{\partial \theta}{\partial t} \right)^2 + \left(\frac{\partial \theta}{\partial x} \right)^2 + \left(\frac{\partial \theta}{\partial y} \right)^2 \right]$$

$$\pm \left[1 - \Delta K(x, y) \right] \sin^2 \theta.$$
(1)

The corresponding equation of motion for the scalar field $\theta(x, y, t)$ has the following form

$$\frac{\partial^2 \theta}{\partial t^2} - \frac{\partial^2 \theta}{\partial x^2} - \frac{\partial^2 \theta}{\partial y^2} + \frac{1}{2} K(x, y) \sin 2\theta = 0,$$
(2)

where the function K(x, y) defines the interaction of the field $\theta(x, y, t)$ with the impurity.

2. One-dimensional case

The sine-Gordon model with the impurity extended in one dimension is defined by the Lagrangian (1) with

$$K(x) = \begin{cases} 1, & x < x_1, \ x > x_2 \\ 1 - \Delta K, \ x_1 \le x \le x_2 \end{cases}$$
(3)

where $W = x_2 - x_1$ is the width of the impurity. Clearly, $\Delta K > 0$ describes a potential well, while $\Delta K < 0$ describes a potential barrier. In the case K(x) = 1 ($\Delta K = 0$) Equation (2) supports the exact solution in the form of topological soliton or, in other words, kink:

$$\theta(\mathbf{x}, t) = 2 \arctan\left(\exp[\gamma(\mathbf{v}_0)(\mathbf{x} - \mathbf{v}_0 t)]\right),\tag{4}$$

where $\gamma(v_0) = (1 - v_0^2)^{-1/2}$ with a parameter $0 < v_0 < 1$ defining the velocity of the kink. Equation (2) also admits the periodic in time solution in the form of a breather,

$$\theta(x,t) = 2 \arctan\left(\frac{\sqrt{1-\omega^2}\sin\omega t}{\omega\cosh\left[\sqrt{1-\omega^2}(x-x_0)\right]}\right),\tag{5}$$

where ω is the breather frequency and x_0 is the coordinate of its center.

The case of $K(x) = 1 - \varepsilon \delta(x)$ where $\delta(x)$ is the Dirac delta function and $0 < \varepsilon < 1$ is a constant has been studied in [5]. It has been demonstrated that in the frame of the underformable kink approximation the impurity acts as a potential and for the chosen sign of ε the potential is attractive and hence the soliton can be localized on the impurity. In the case of deformable kink, in addition to the oscillatory motion of the kink in the potential generated by the impurity, a strong modification of the kink shape, having a resonant character, can take place. The possibility of excitation of the impurity mode as a result of kink scattering that results in a considerable change of kink dynamics has also been considered. For the case of finite size impurity of the form (3) the interaction of the kink with the impurity has been analyzed for both non-deformable and deformable kinks [20, 22, 24].

Let us investigate the relation between the spectra of the impurity mode and the small-amplitude excitations of (1). We take into account that for the one-dimensional case of (2) with K(x) = 1 there exists the vacuum solution $\theta_{\pm}(x, t) = 0$. We look for the spectrum of the small-amplitude vibrations in the vicinity of this solution

$$\theta(\mathbf{x},t) = \theta_{\pm} + \delta\theta(\mathbf{x},t), \quad \delta\theta(\mathbf{x},t) \ll 1.$$
 (6)

Substituting (6) into (2) after linearization with respect to $\delta \theta$, one gets the equation

$$L\delta\theta(\mathbf{x}) = \omega_n^2 \delta\theta(\mathbf{x}),\tag{7}$$

which is the Schrödinger equation with the operator $L = -\frac{d^2}{dx^2} + K(x)$, where ω_n is the impurity mode frequency. Let us look for the localized solutions of the Schrödinger equation (7). It is convenient to introduce the following notations

$$\chi^2 = 1 - \omega^2, \quad k^2 = \omega^2 - (1 - \Delta K).$$
 (8)

For K(x) defined by (3) even and odd solutions to (7) are possible [35]

$$\Psi_{+} = \begin{cases} A_{1}^{+} e^{\chi x}, & x < -W/2 \\ B_{2}^{+} \cos kx, & -W/2 \le x \le W/2 \\ B_{3}^{+} e^{\chi x}, & x > W/2 \end{cases}$$
(9)

$$\Psi_{-} = \begin{cases} A_{1}^{-} e^{\chi x}, & x < -W/2 \\ B_{2}^{-} \sin kx, & -W/2 \le x \le W/2 \\ B_{3}^{-} e^{\chi x}, & x > W/2 \end{cases}$$
(10)

We subject the solution to the condition of smoothness and continuity at the point x = W/2 and obtain the following dispersion relations for even (9) and odd (10) solutions, respectively,

$$\tan\left(k\frac{W}{2}\right) = \frac{\chi}{k},\tag{11}$$

$$\tan\left(k\frac{W}{2}\right) = -\frac{\chi}{k}.$$
(12)

Because, in the considered case, ΔK and W are constant, equations (11,12) give the possibility to find all frequencies ω of the impurity modes in the potential well of given size. The states with even and odd wave functions alternate and the first odd solution, that corresponds to the second localized state, appears when the relation $\pi^2 = W^2 \Delta K$ is satisfied.

Let us study numerically the large-amplitude localized impurity modes excited due to the interaction with a kink for the case $\Delta K \leq 0$. The most interesting case is when the size of the soliton is of the same order with the size of the impurity because in this case the soliton shape is strongly affected by the impurity. To solve the equations of motion (2) numerically we use the iteration method for the explicit scheme. The following algorithm was applied. Initially we have a SGE kink (4) moving with a constant speed. Boundary conditions have the form $\theta(\pm \infty) = 0$, π ; $\theta'(\pm \infty) = 0$. We introduce the mesh for the spatial coordinate and iterate with respect to time, taking into account the convergence condition for the explicit scheme, to find the kink position at the next time step. The characteristics of the nonlinear wave were found from the numerically constructed function $\theta(x, t)$.

The numerical experiments have demonstrated that the kink passing the impurity excites the bell-shaped nonlinear oscillatory wave. We found that this oscillatory mode can be well fitted by the expression

$$\theta^*(x,t) = A \exp(-\alpha(t-t_0)) \arctan\left(\frac{\sqrt{1-\omega^2}\sin[\omega\gamma(t-t_0)]}{\omega\cosh\left(\gamma\sqrt{1-\omega^2}x\right)}\right),\tag{13}$$

which is a stationary breather (5) in the presence of damping with the coefficient α and with an additional fitting parameter γ . We compare the numerically found time evolution of the field value at the center of the impurity (x = 0) with that found from the approximating formula (13). The numerical result and the result obtained from the fitting formula (13) overlap. Damping of the breather is due to slow radiation of small-amplitude extended waves. Amplitude of the breather excited by the passing kink depends on the kink velocity v_0 (Figure 1) and the curve has a maximum with the value dependent on the impurity parameters ΔK and W.

The amplitude of the excited breather also depends on ΔK and W and it vanishes when $\Delta K \rightarrow 0$, $W \rightarrow 0$, i.e., in the absence of impurity.

Breather frequency ω_B is practically independent of the kink velocity v_0 , while it depends on ΔK and W as shown in Figure 2.

It can be seen that for $\Delta K \rightarrow 0$, $W \rightarrow 0$ the breather frequency tends to unity but the frequencies of kink's translational and pulsation (oscillation of kink wigth) modes excited due to the interaction with impurity tend to zero [20]. This behavior can be easily understood



Figure 1. Breather amplitude A_{max} measured at the impurity center as the function of kink velocity v_0 for (a) W = 1 and (b) W = 1.5. Curves 1 to 6 are for $\Delta K = \{0.5, 0.75, 1, 1.25, 1.5, 1.75\}$, respectively.





taking into account that breather energy scales as $E \sim (1-\omega_B^2)^{1/2}$ [2] meaning that for vanishing size of the impurity the energy and the amplitude of the breather also vanish. It can also be seen that the linear approximation for the first even solution described above gives a good description of the excited breathers in the range of small amplitudes because the frequencies calculated from the analytical expression (11) (solid lines in Figure 2) coincide with the frequencies found numerically by solving (2). Excitation of the first odd mode (the second main state) will be discussed for the case of kink pinning by the impurity. In this case, as it was already mentionaed, pulsation and translational modes can be excited on the kink [20]. In the case when kink pulsation mode frequency $\omega_{pulse} < 1$, after a transient period a state is formed that can be described with a good accuracy by the kink-type solution. Therefore, in our case the most interesting is the range of parameters ΔK and W where $\omega_{pulse} \rightarrow 1$ and the excited nonlinear wave (Figure 3(a)) differs considerably from the kink solution (Figure 3(b)). Algebraic difference of the solutions has the form dramatically different from 13 and it is analogous to the form of the odd solutions to the Schrödinger equation.

One should also take into account that for the case of large-amplitude nonlinear waves obtained numerically, a three-kink solution can be not just a linear sum of the kink and breather solutions but a modified solution that takes into account kink vibrations described by the wobble SGE solution [36, 37]. For yet increasing values of ΔK and W let us take into account that, as it has been shown in [4], at the critical value of parameters,

$$W(1 - \Delta K) = 2, \tag{14}$$

in the vicinity of the impurity a stable static soliton can exist whose amplitude can be estimated as

$$\cos A_{\max} = 2/[W(1 - \Delta K)]. \tag{15}$$

Numerical results show that for sufficiently large ΔK and W after the passing of kink, in the vicinity of the impurity a soliton is formed. The dependence of soliton amplitude on ΔK and W (Figure 4) can be approximately expressed as $\cos A = 1.8/[W(1 - \Delta K)]$. In Figure 5 the ranges of the impurity parameters where the breather and the soliton can exist are presented and for comparison the curve defined by (14) is plotted.



Figure 3. (a) Wobble kink profile and (b) its difference from the kink solution at different times, 1 for t = 1179.37, 2 for t = 1180.87, and 3 for t = 1182.37, for the case W = 1, $\Delta K = 1.2$ and initial kink velocity $v_0 = 0.2$. The impurity center is at $x^* = -0.5$.



Figure 4. The dependence of the stabilized soliton amplitude on the impurity width *W* at fixed $\Delta K = 2.8$. Curve 1 gives numerical result, while curve 2 corresponds to (15).



Figure 5. Space of impurity parameters showing the range where passing kink excites a breather (above the lines 1 and 2), or a soliton (below the lines 1 and 2). Line 1 was found numerically and line 2 is given by (14).

For definiteness let us take the impurity in the following form [18]

$$K(x,y) = \begin{cases} 1, & x < x_1, x > x_2, y < y_1, y > y_2, \\ 1 - \Delta K, & x_1 \le x \le x_2, y_1 \le y \le y_2, \end{cases}$$
(16)

where $W_x = x_1 - x_2$ and $W_y = y_1 - y_2$ specify the size of the impurity. Similarly to the one-dimensional case, Equation (2) was integrated with the use of an explicit scheme. A uniform mesh with the step ξ was introduced for x and y coordinates,

$$x_i = \xi i, \ i = -N_x, ..., N_x, \quad y_i = \xi j, \ j = -N_y, ..., N_y,$$
(17)

and for the time t the mesh with the time step τ was employed,

$$t_n = \tau n, \ n = 0, 1, ..., N_t,$$
 (18)

where N_x , N_y , N_t are integers. For N_x and N_y we took the values from 512 to 2048. It was confirmed that the number of mesh points for the spatial coordinates does not noticeably affect the numerical results. Initially we have a SGE kink (4) moving with constant velocity. The boundary conditions for the *x*-coordinate have the form $\theta(\pm N_x\xi, y) = \theta_0(\pm N_x\xi)$; $\theta'(\pm N_x\xi, y) = \theta'_0(\pm N_x\xi)$ and for the *y*-coordinate free edges are simulated.

Let us consider the case of $\Delta K > 0$ when, as it was shown above, nonlinear waves localized at the impurity are excited due to the interaction with the kink. In Figure 6 the process of kink-impurity interaction is presented and one can see that the nonlinear localized wave radiating extended waves is excited. The wave is called here breathing pulson. If $W_x = W_y$ then the breathing pulson is symmetric with respect to x and y axis (Figure 7(a) and (b)) and the symmetry is lost for $W_x \neq W_y$ (Figure 7(c) and (d)). Firstly, the excited localized wave has a bell shape and later (t > 20, curve 1 in Figure 8) the periodic oscillations can be seen at the center of the impurity having coordinates (x^*, y^*). With increase in the impurity size the oscillation frequency reduces. The amplitude decreases with time owing to the radiation of extended waves. It should be noted that the extended waves cannot be described by the harmonic function because they have a nonlinear nature (Figure 9). The breathing pulson can be regarded as a long-lived nonlinear excitation.



Figure 6. The excitation and evolution of a breathing pulson for the case $W_x = 1$, $W_y = 3$, and $\Delta K = 2$. (a) t = 11.55, (b) t = 12.6, (c) t = 13.65, (d) t = 15.75, (e) t = 17.43, and (f) t = 19.32. The impurity center coordinates are $x^* = y^* = 60$.



Figure 7. Snapshots of the functions $\theta(x, y^*, t)$ (a,c) and $\theta(x^*, y, t)$ (b,d) for the case $\Delta K = 2$, $v_0 = 0.57$. In (a,b) the impurity is symmetric, $W_x = W_y = 1$, and the curves 1 to 6 are for $t = \{28.2, 29.04, 29.5, 30.24, 30.72, 31.5\}$. In (c,d) the impurity is asymmetric, $W_x = 1$, $W_y = 3$, and the curves 1 to 6 are for $t = \{44.28, 45.6, 45.96, 46.5, 46.86, 48.18\}$. The impurity center coordinates are $x^* = y^* = 60$.

Copyright © 2016 John Wiley & Sons, Ltd.



Figure 8. The function $\theta(x^*, y^*, t)$ for the case $v_0 = 0.85$, $W_x = 1$, $W_y = 3$, $\Delta K = 2$ (curve 1) and $\Delta K = 5$ (curve 2).



Figure 9. Snapshots of $\theta(x, y^*, t)$ (a) and $\theta(x^*, y, t)$ (b) for $W_x = 1$, $W_y = 1$, $\Delta K = 2$. Curves 1 to 5 correspond to $t = \{42, 54, 66, 78, 90\}$.



Figure 10. The dependence of the breathing pulson oscillation frequency ω_p (a) on the parameter ΔK for the case $W_y = 3$ and $W_x = 1$ (curve 1), $W_x = 2$ (curve 2), and $W_x = 3$ (curve 3); (b) on the parameter W_x for the case $W_y = 1$ and $\Delta K = 2$ (curve 1), $\Delta K = 3$ (curve 2), and $\Delta K = 4$ (curve 3). Kink initial velocity is $v_0 = 0.57$.

It is well-known that SGE supports the 2D solution called pulson describing long-lived spatially localized vibrations [28–32]. The numerically found breathing pulson can be well approximated by the expression

$$\theta(r,t) = A \arctan\left[\frac{\sqrt{1-\omega_{\rho}^{2}}}{\omega_{\rho}}\operatorname{sech}\left(\sqrt{1-\omega_{\rho}^{2}}Br\right)\sin[\omega_{\rho}(t-t_{0})]\right],$$
(19)

where $r = \sqrt{(x/\Delta_x)^2 + (y/\Delta_y)^2}$, with Δ_x and Δ_y being the pulson width along x and y axis, respectively. For example, for the curve 1 in Figure 8 one should set in (19) r = 0 and A = 0.42, $\omega_P = 0.85$, B = 4.8, $t_0 = 2.5$, $\Delta_x = 2.4$, and $\Delta_y = 3.4$.

Our numerical results show that the breathing pulson frequency ω_P [determined from $\theta(x^*, y^*, t)$] does not depend on kink velocity υ_0 , but it is a function of W_x , W_y and ΔK . In Figure 10 the dependence of ω_P on the parameters ΔK and W_y is shown. It can be seen that with decrease in the impurity size the breathing pulson frequency (similar to what was observed for the breather) tends to unity. Dependence of ω_P on $K = 1 - \Delta K$ can be approximately given by

$$\omega_P = (a(|K^*| + K))^q / \sqrt{1 + (a(|K^*| + K))^{2q}},$$
(20)

where *a* is a constant, $q \approx 2$, K^* is the smallest value of *K* when breathing pulson is still formed in the vicinity of the impurity. Dependence of ω_P on W_y (as well as on W_x) can be approximately expressed as $\omega_P = 1 - (bW_y)^P / \sqrt{1 + (bW_y)^{2p}}$, where *b* is a constant and $p \approx 2$. Maximal breathing pulson amplitude, A_{max} , as the function of the kink velocity v_0 is presented in Figure 11 and, similarly to the one-dimensional case, it has a maximum. With decrease in the impurity size A_{max} vanishes. The dependence $A_{max}(\Delta K)$ for small values of ΔK is close to linear. The dependence $A_{max}(W_y)$ for $W_y < 1$ is close to linear and for large W_y it saturates. Note that for fixed v_0 maximal value of A_{max} strongly depends on ΔK , W_x and W_y .

For increasing values of the parameters ΔK , W_x and W_y , after the kink passes the impurity, a localized nonlinear wave called here breathing 2D soliton is excited on the impurity (Figure 12). The breathing 2D soliton shape depends on the parameters W_x and W_y and it can be symmetric (Figure 13(a) and (b)) or asymmetric (Figure 13(c) and (d)). Our numerical results suggest that the breathing



Figure 11. The dependence of the breathing pulson maximum amplitude A_{max} at the impurity center on the kink velocity v_0 for the case $W_x = 1$, $\Delta K = 2$, and $W_y = 1$ (curve 1), $W_y = 2$ (curve 2), $W_y = 3$ (curve 3).



Figure 12. Excitation and evolution of the breathing 2D soliton for the case $W_x = 1$, $W_y = 3$, $\Delta K = 5$. Panels (a) to (f) correspond to $t = \{8, 9.4, 11, 14, 20, 26\}$. The impurity center coordinates are $x^* = y^* = 60$.



Figure 13. Snapshots of the functions $\theta(x, y^*, t)$ (a,c) and $\theta(x^*, y, t)$ (b,d) for the case (a,b) $W_x = 2$, $W_y = 2$, and t = 30.18 (curve 1), t = 31.8 (curve 2), t = 33.72 (curve3); (c,d) $W_x = 2$, $W_y = 3$, and t = 30.78 (curve 1), t = 32.4 (curve 2), t = 34.02 (curve 3). $\Delta K = 5$, $v_0 = 0.85$. The impurity center coordinates are $x^* = y^* = 60$.

2D soliton is a long-lived excitation with amplitude slowly decreasing in time. The breathing 2D soliton cannot be described by the direct sum of 2D soliton solution and pulson solution to SGE. On the other hand, the breathing 2D soliton can be well approximated by the expression

$$\theta(r,t) = \arctan\left(\frac{A_0 + A_1 \sin[\omega_5(t-t_0)]}{\omega_5^{-1}\sqrt{1-\omega_5^2}\cosh(\sqrt{1-\omega_5^2}Br)}\right),$$
(21)

where $r = \sqrt{(x/\Delta_x)^2 + (y/\Delta_y)^2}$, $B = B_0 - B_1 \sin[\omega_5(t - t_0)]$, ω_5 is breathing 2D soliton frequency, A_0 , A_1 , B_0 , B_1 are the breathing 2D soliton parameters that depend on the impurity parameters, Δ_x , Δ_y are the characteristic widths of the soliton along X and Y axis, respectively. For example, the curve 2 in Figure 8 is well fitted by (21) with r = 0 and $A_0 = 0.63$, $A_1 = 0.13$, $B_0 = 9$, $B_1 = 1$, $\omega_5 = 0.94$, $t_0 = 0$, $\Delta_x = 1.8$, and $\Delta_y = 3.2$.

Our numerical results have demonstrated that the breathing 2D soliton frequency, ω_s , similarly to what was observed for the pulson, does not depend on the initial kink velocity but is a function of the parameters W_x , W_y and ΔK (Figure 14). It can be seen that



Figure 14. The dependence of the breathing 2D soliton oscillation frequency ω_s (a) on the parameter ΔK for the case $W_y = 2$ and $W_x = 2$ (curve 1), $W_x = 3$ (curve 2), $W_x = 4$ (curve 3); and (b) on the parameter W_y for the case $W_x = 1$ and $\Delta K = 5$ (curve 1), $\Delta K = 5.5$ (curve 2), $\Delta K = 6$ (curve 3). Initial kink velocity is $v_0 = 0.85$.



Figure 15. The breathing 2D soliton width Δ_y as the function of the parameter W_y for $\Delta K = 5$, $v_0 = 0.85$, and $W_x = 2$ (curve 1), $W_x = 3$ (curve 2), $W_x = 4$ (curve 3).



Figure 16. The regions of the impurity parameters allowing the existence of the breathing pulson (above the lines 1, 2 and 3), and the breathing 2D soliton (below the lines 1, 2 and 3). The lines 1 and 2 present numerical results for $W_y = 1$ and $W_y = 3$, respectively, while line 3 is plotted with the help of (14).

 $\omega_{\rm S}$ tends to unity for increasing W_x , W_y and ΔK . The dependence of $\omega_{\rm S}$ on $K = 1 - \Delta K$ can be approximated by the expression $\omega_{\rm S} = (c(|K^*| - K))^q / \sqrt{1 + (c(|K^*| - K))^{2q}}$, where $q \approx 6$, c is a constant, and K^* is the largest value of K at which the breathing 2D soliton can be formed in the impurity region. The dependence of $\omega_{\rm S}$ on W_y (as well as on W_x) can be approximately given by $\omega_{\rm S} = (aW_y)^p / \sqrt{1 + (aW_y)^{2p}}$, where $p \approx 3$, and a is a constant. In Figure 15 for the breathing 2D soliton we present the dependence of Δ_y on the parameters W_y . It can be seen that Δ_y depends almost linearly on W_y . The relation between parameters describing the size of impurity and the width of the breathing 2D soliton can be approximately written as

$$\frac{\Delta_x^2}{W_x^2} + \frac{\Delta_y^2}{W_y^2} = R^2, \tag{22}$$

where for the symmetric impurity region $R = 2^{3/4}$, and, in general, parameter R depends on the parameters ΔK , W_x and W_y .

In Figure 16 we plot the regions of the parameters ΔK , W_x and W_y where different localized nonlinear excitations exist. It can be seen that the increase in W_y shifts the critical curves toward the smaller values of the parameters ΔK and W_x .

4. Conclusions

Using analytical and numerical methods the dynamics of the sine-Gordon equation kinks passing through the attractive impurity was examined. The cases of extended one-dimensional and two-dimensional impurities were studied.

By linearizing the sine-Gordon equation the dispersion relations for the small-amplitude localized impurity modes were obtained. Using numerical methods the possibility of excitation by the passing kink of the first even and odd modes of high-amplitude localized on the impurity modes was demonstrated. The numerically obtained dispersion relations in the case of low oscillation amplitudes are in a good agreement with the results of analytical calculations.

For the case of two-dimensional impurity, the possibility of excitation by the passing kink of non-linear high-amplitude waves of new type, called here breathing pulson and breathing 2D soliton, was shown numerically. Analytical expressions to model these new excitations were suggested. The analytical expressions are modified exact solutions to the unperturbed 2D SGE to take into account damping due to the slow radiation of extended waves. The breathing pulson and breathing 2D soliton are long-lived and can be of both symmetric and asymmetric type depending on the impurity type. The range of the impurity parameters where the breathing pulson and breathing 2D soliton can be excited was determined. The amplitude, size and frequency of the excited localized nonlinear modes as the functions of the impurity parameters were given.

Our study contributes to the understanding of the interaction of the SGE kink with the inhomogeneities in 1D and 2D cases.

References

- 1. Christiansen PL, Sorensen MP, Scott AC (eds). Nonlinear science at the dawn of the 21st century. Springer: Berlin, 2000.
- 2. Scott AC (ed). Encyclopedia of Nonlinear Science. Routledge: New York, 2004.
- 3. Yakushevich LV, Savin AV, Manevitzch LI. On the internal dynamics of topological solitons in DNA. Physical Review E 2002; 66:016614–016629.
- 4. Ekomasov EG, Azamatov SA, Murtazin RR. Studying the nucleation and evolution of magnetic inhomogeneities of the soliton and breather type in magnetic materials with local inhomogeneities of anisotropy. *The Physics of Metals and Metallography* 2008; **105**(4):313–321.
- 5. Braun OM, Kivshar YS. The Frenkel-Kontorova Model: Concepts, Methods, and Applications. Springer-Verlag: Berlin, 2004.
- 6. Cuevas-Maraver J, Kevrekidis PG, Williams F (eds). The Sine-Gordon Model and Its Applications: From Pendula and Josephson Junctions to Gravity and High-energy Physics, Vol. 10. Springer, 2014.
- 7. Fogel MB, Trullinger SE, Bishop AR, Krumhandl JA. Dynamics of sine-gordon solitons in the presence of perturbations. *Physical Review B* 1977; **15**:1578–1592.
- 8. Kivshar YS, Pelinovsky DE, Cretegny T, Peyrard M. Internal modes of solitary waves. *Physical Review Letters* 1998; **80**:5032–5035.
- 9. Kivshar YS, Malomed BA, Zhang F, Vazquez L. Creation of sine-gordon solitons by a pulse force. Physical Review B 1991; 43:1098–1109.
- 10. Ekomasov EG, Gumerov AM, Kudryavtsev RV. On the possibility of the observation of the resonance interaction between kinks of the sine gordon equation and localized waves in real physical systems. *JETP Letters* 2015; **101**(12):835–839.
- 11. Ekomasov EG, Gumerov AM, Murtazin RR, Kudryavtsev RV, Ekomasov AE, Abakumova NN. Resonant dynamics of the domain walls in multilayer ferromagnetic structure. Solid state phenomena 2015; 233-234:51–54.
- 12. Ekomasov EG, Murtazin RR, Bogomazova OB, Gumerov AM. One-dimensional dynamics of domain walls in two-layer ferromagnet structure with different parameters of magnetic anisotropy and exchange. *Journal of Magnetism and Magnetic Materials* 2013; **339**:133.
- 13. Dmitriev SV, Kevrekidis PG, Kivshar YS. Radiationless energy exchange in three-soliton collisions. *Physical Review E* 2008; **4**(78):046604.
- 14. Gumerov AM, Ekomasov EG, Murtzin RR, Nazarov VN. Transformation of sine-gordon solitons in models with variable coefficients and damping. Computational Mathematics and Mathematical Physics 2015; **55**(4):628–637.
- 15. González JA, Bellorin A, Guerrero LE. Internal modes of sine-gordon solitons in the presence of spatiotemporal perturbations. *Physical Review E (Rapid Communications)* 2002; **65**:065601.
- 16. Chacón R, Bellorín A, Guerrero LE, González JA. Spatiotemporal chaos in sine-gordon systems subjected to wave fields: Onset and suppression. *Physical Review E* 2008; **77**(4):046212.
- 17. Willis CR. Comment on existence of internal modes of sine-gordon kinks. Physical Review E 2006; 73:068601.
- 18. Bratsos AG. The solution of the two-dimensional sine-gordon equation using the method of lines. *Journal of Computational and Applied Mathematics* 2007; **206**(1):251–277.
- 19. Currie JP, Trullinger SE, Bishop AR, Krumhandl JA. Numerical simulation of sine-gordon soliton dynamics in the presence of perturbations. *Physical Review B* 1977; **15**(12):5567–5580.
- 20. Ekomasov EG, Shabalin MA. Simulation the nonlinear dynamics of domain walls in weak ferromagnets. *The Physics of Metals and Metallography* 2006; **101**:S48–S50.
- 21. Gumerov AM, Ekomasov EG, Zakiryanov FK, Kudryavtsev RV. Structure and properties of four-kink multisolitons of the sine-gordon equation. *Computational Mathematics and Mathematical Physics* 2014; **54**(3):491–504.
- 22. Paul DI. Soliton theory and the dynamics of a ferromagnetic domain wall. Journal of Physics C: Solid State Physics 1979; 12:585–593.
- 23. Zhang F, Kivshar YS, Vazquez L. Resonant kink-impurity interactions in the sine-gordon model. *Physical Review A* 1992; **45**(8):6019–6030.
- 24. Piette B, Zakrzewski W J. Scattering of sine-gordon kinks on potential wells. Journal of Physics A: Mathematical and Theoretical 2007; 40:5995–6010.
- 25. Piette B, Zakrzewski WJ, Brand J. Scattering of topological solitons on holes and barriers. *Journal of Physics A: Mathematical and General* 2005; **38**(48):10403–10412.
- 26. Piette B, Zakrzewski WJ. Dynamical properties of a soliton in a potential well. Journal of Physics A: Mathematical and Theoretical 2007; 40(2):329–346.
- 27. Javidan K. Analytical formulation for soliton-potential dynamics. *Physical Review E* 2008; **78**:046607.
- 28. Dodd RK, Eilbeck JC, Gibbon JD, Morris HC. Solitons and nonlinear wave equations. Academic Press: London, 1982.
- 29. Johnson S, Suarez P, Biswas A. New exact solutions for the sine-gordon equation in 2+1 dimensions. *Computational Mathematics and Mathematical Physics* 2012; **52**(1):98–104.
- 30. Wazwaz AM. N-soliton solutions for the sine-gordon equation of different dimensions. *Journal of Applied Mathematics and Informatics* 2012; **30**(5-6):925–934.
- 31. Qing M, Bin H, Weiguo R, Yao L. New exact solutions of the (n+1)-dimensional sine-gordon equation using double elliptic equation method. International Journal of Computer Mathematics 2010; 87(3):591–606.
- 32. Koutvitsky VA, Maslov EM. Parametric instability of the real scalar pulsons. *Physics Letters A* 2005; **336**(1):31–36.
- 33. Dehghan M, Shokri A. A numerical method for solution of the two-dimensional sine-gordon equation using the radial basis functions. *Mathematics and Computers in Simulation* 2008; **79**(3):700–715.
- 34. Ekomasov EG, Azamatov SA, Murtazin RR. Excitation of nonlinear solitary flexural waves in a moving domain wall. *The Physics of Metals and Metallography* 2009; **108**(6):532–537.
- 35. Landau LD, Lifshitz EM. Quantum Mechanics: Non-Relativistic Theory, Vol. 3. Butterworth-Heinemann, 1977.
- 36. Ferreira LA, Piette B, Zakrzewski WJ. Wobbles and other kink-breather solutions of sine-gordon model. *Physical Review E* 2008; **77**:036616.
- 37. Kalberman G. The sine-gordon wobble. Journal of Physics A: Mathematical and General 2004; 37:11603–11612.